


## AMENDMENTS TO THE CLAIMS

This listing of the claims will replace all prior versions including the claims in the application.

Listing of the Claims:

### CLAIMS:

1. (Currently Amended) A benzotriazole of formula I,



I

in which:

n is an integer selected from 0, 1 and 2

R1 to R8 are each H, except that

one of R2 and R3 may be selected from

Br, Cl, CH<sub>3</sub>, CN, NH<sub>2</sub>, NO<sub>2</sub>, CF<sub>3</sub>, OCH<sub>3</sub>, phenoxy, benzoyl, CH(OH)-phenyl, S-cyclohexyl, and CO-OCH<sub>3</sub>; or

R1 is Cl and R3 is CF<sub>3</sub>; or

R2 is F and R3 is Cl; or

one of R6 and R7 may be selected as follows:


R6 is CH<sub>3</sub> or R7 is selected from CH<sub>3</sub>, C<sub>2</sub>H<sub>5</sub>, CH(CH<sub>3</sub>)<sub>2</sub>, C(CH<sub>3</sub>)<sub>3</sub>, CF<sub>3</sub>, Br, Cl, benzyl and CO-OC<sub>2</sub>H<sub>5</sub>; or

R6 and R7 each are CH<sub>3</sub>; or

R6 and R7 may be replaced by a double bond between the ring carbon to which they are attached; or R5 and R6 or R6 and R7 may, in combination with the carbon to which they are attached, form a benzo-fused ring or, when n is 0, may form cyclohexanediyI, the ring formed from R6, R7 and the carbons to which they are attached being optionally substituted singly by NH<sub>2</sub> or NO<sub>2</sub> or substituted singly or doubly by OCH<sub>3</sub>; and

R7 and R8, together with the carbon to which they are attached, may form cyclopentyl, diazirinediazirine or =CH<sub>2</sub>; provided, however, that when R1 to R5 and R8 are H, and R6, R7 and the carbons to which they are attached form a benzo-fused ring, n is not 1, and when R1 and R3-R8 are H and R2 is CH<sub>3</sub>, n is not 1 and when R1 to R8 are each H, n is 2.

2. (Original) A benzotriazole of formula I as claimed in claim 1,



in which:

R1 to R8 are each H, except that one of R2 and R3 may instead be selected from the following substituents:

R2 is selected from Br, Cl, CN, NO<sub>2</sub>, CF<sub>3</sub>, OCH<sub>3</sub>, phenoxy, benzoyl, CH(OH)-phenyl, S-cyclohexyl, and CO-OCH<sub>3</sub>; or

R3 is selected from CH<sub>3</sub>, CN, Br, Cl, NH<sub>2</sub>, NO<sub>2</sub>, and benzoyl.

3. (Original) A benzotriazole of formula I as claimed in claim 1, in which :

R1 to R8 are each H; except that one of R2 and R3 may instead be selected from the following substituents:

R2 is selected from Br, Cl, NO<sub>2</sub>, OCH<sub>3</sub>, phenoxy, and CO-OCH<sub>3</sub>; or

R3 is NH<sub>2</sub>; or

R2 is F and R3 is Cl; or

n is 1 or 2 and

one of R6 and R7 is selected from the following substituents:

R6 is CH<sub>3</sub>; or R7 is selected from CH<sub>3</sub>, CF<sub>3</sub> and Br; or

R6 and R7 may be replaced by a double bond between the ring carbons to which they are attached; or

R6 and R7 may, in combination with the carbons to which they are attached, form a benzo-fused ring, which may be optionally substituted singly by NH<sub>2</sub> or substituted singly or doubly by OCH<sub>3</sub>; and

R7 and R8 together, with the carbon to which they are attached, form a cyclopentyl; or

n is 0 and

R6 and R7, together with the carbons to which they are attached, form a benzo-fused ring or cyclohexanediyi.

4. (Original) A benzotriazole of formula I as claimed in claim 1, wherein R1 to R8 are each H; except that one of R2 and R3 may instead be selected from the following substituents:

R2 is selected from Br, CN, CF<sub>3</sub>, OCH<sub>3</sub>, phenoxy, benzoyl, CH(OH)-phenyl, and S-cyclohexyl; or

R3 is selected from CN, Br, Cl, NO<sub>2</sub>, and benzoyl; or

R1 is Cl and R3 is CF<sub>3</sub>; or

n is 1 and

one of R6 and R7 is selected from the following substituents:

R6 is CH<sub>3</sub>; or



R7 is selected from CH<sub>3</sub>, C<sub>2</sub>H<sub>5</sub>; CH(CH<sub>3</sub>)<sub>2</sub>, C(CH<sub>3</sub>)<sub>3</sub>, benzyl and CO-OC<sub>2</sub>H<sub>5</sub>; or







R6 and R7 are each CH<sub>3</sub>; or R6 and R7 may be replaced by a double bond between the ring carbons to which they are attached; or





R5 and R6 or R6 and R7 may, in combination with the carbons to which they are attached, form a benzo-fused ring;

provided, however, that when R1 to R5 and R8 are H and R6, R7 and the carbon to which they are attached form a benzo-fused ring, n is not 1.



5. (Currently Amended) A benzotriazole ~~of claim 1~~ selected from the group consisting of the compounds having the following structures:














,









۳






6. (Currently Amended) A benzotriazole of ~~claim 1~~ selected from the group consisting of the compounds having the following structures:







3




1









1















3






3













, and




7. (Original) A benzotriazole as claimed in claim 6 selected from the group consisting of the compounds having the following structures:







,




,




,







,




,








,




, and



8. (Original) A benzotriazole as claimed in claim 7 selected from the group consisting of the compounds having the following structures:





and



9. (non-elected) A process for preparing the compounds of formula I as claimed in claim 1, which comprises

- a) acylating benzotriazole 2 with carbamoyl chlorides 3, or
- b) initially reacting benzotriazoles 2 with phosgene and then reacting the resulting benzotriazolecarbonyl chlorides 5 with amines or anilines to give the compounds of the formula I, in which the substituents are as defined in claim 1:



10. (non-elected) A medicament comprising, as an active ingredient, a benzotriazole of formula I as claimed in claim 1.

11. (non-elected) A method of inhibiting hormone-sensitive lipase, HSL, comprising administering to a patient in need thereof an effective amount of a medicament as claimed in claim 10.
12. (non-elected) A method of treatment of non-insulin-dependent diabetes mellitus or of diabetic syndrome or syndrome X, comprising administering to a patient in need thereof an effective amount of a medicament as claimed in claim 10.
13. (non-elected) A method for the treatment of non-insulin-dependent diabetes mellitus or of diabetic syndrome X, comprising administering to a patient in need thereof an effective amount of at least one benzotriazole of formula I as claimed in claim 1.